

STATISTICAL ANALYSIS PLAN ENDOSCRATCH TRIAL NCT03108157

Study title:

Endometrial Scratch Effect on Pregnancy Rates in Patients Undergoing Egg-donation IVF

ClinicalTrials.gov ID:

NCT03108157

Version

25th July 2019

Protocol Version

2.0 2nd November 2016 (1.PROTOCOLO SCRATCH V 2.0.docx.pdf)

Data Collection Form Version

1.0 2nd November 2016 (2. CRD.pdf)

Signatures

Principal investigator	SAP author	SAP reviewer
Alexandra Izquierdo	David Lora	Alexandra Izquierdo

Note: Add name, surname and date next to the signature.

SAP reviews

Protocol Version	SAP review	Section modified	Description and reason for modification	Date of modification

1 Introduction

1.1 Background

Embryo implantation remains one of the main challenges in assisted reproduction. Relevant improvements have been accomplished in reproductive medicine: different protocols for controlled ovarian stimulation and endometrial preparation, embryo culture with time-lapse technologies, embryo pre-implantational genetic testing and endometrial genetic assessment for implantation potential. Despite the fact that these changes have led to increasing pregnancy rates in the last few years, the implantation process is still inefficient, as it remains around 30% of all embryos replaced (1), and it is not yet totally understood.

1.2 Study justification

Several studies have tried to determine whether an endometrial injury performed in the cycle preceding the embryo transfer in in vitro fertilization (IVF) cycles could enhance embryo implantation. Barash et al (2) reported for the first time a two-fold increase in pregnancy rate in patients that had undergone multiple ES before the IVF cycle, compared to those patients who had no ES performed.

Since then, many authors have tried to determine ES effects after controlled ovarian stimulation (COS), but while some of them have found an increase in pregnancy rates (2–5), many others have been unable to find such differences (6–11). The main limitation in reaching a conclusion is that most of these are underpowered observational studies, with a low number of patients included, with differences in timing (luteal or follicular phase from the preceding or same cycle), number of ES (one, two or more procedures), type of catheter and different stimulation protocols. It is important to note that those studies that have found some positive effects of ES have included patients with implantation failures (5,12) whereas those that included patients in their first or second IVF cycle were unable to find any differences (8,9). It is also relevant that some studies included as control patients, those who had undergone a hysteroscopy prior to the IVF cycle and, even if an ES was not performed in these patients, we may assume that the endometrium was exposed to some “damage” as well (13). Another study included a cervical biopsy for those patients included in the control group, what cannot be really considered as “placebo” (10).

A systematic review conducted by Potdar et al (14) including 7 studies with 2062 patients, found a three-fold increase in pregnancy rates in those patients that received ES. Similar results were also found some time later by a Cochrane Review by Nastri et al (15) with moderate-quality evidence, signaling the need for well-designed trials without uterine instrumentation in the control group, stratification for implantation failure and the necessity to report live birth rates. This review also showed that endometrial injury on the day of oocyte retrieval decreased live birth (RR 0.31, 95% CI 0.14 to 0.69) and clinical pregnancy (RR 0.36, 95% CI 0.18 to 0.71).

All these studies were conducted after COS, but there is only one retrospective study in patients receiving embryos from donor eggs, and who have not undergone ovarian stimulation (14). When comparing egg donation cycles to other IVF treatments, we find two main differences: the first one is that embryo quality is presumably optimal, since all embryos come from donor eggs, avoiding the confusion factor of embryo quality according to maternal issues (age, BMI, polycystic ovaries, low ovarian reserve...) and the second difference is that all patients receive hormone replacement therapy with a homogeneous preparation of the endometrium, avoiding different hormonal environments caused by diverse responses to controlled ovarian stimulation in IVF.

1.3 Implantation Failure

Embryo implantation is a complex process that requires successive coordinated steps to allow the embryo attach and invade the inner layers of the endometrium.

Implantation failure definition is very heterogeneous. Coughlan et al (17) considered implantation failure if 4 or more good quality embryos had been replaced in 2 or more embryo transfers.

When there is a suspicion of implantation failure, it is necessary to perform certain tests in order to exclude possible causes: embryo genetic testing, maternal coagulation and immune problems and window of implantation determination and thus endometrial receptivity to the embryo.

The endometrium is a dynamic tissue with a complex architecture that undergoes several changes during the menstrual cycle which have an important impact on embryo implantation. Endometrial scratching (ES) is a simple procedure aiming to create a mild

endometrial injury that has been proposed to improve the embryo-endometrium dialogue. Different authors have attributed this improvement to the effects of different cytokines and growth factors involved in an acute endometrial inflammatory process (18), the enhancement of new vascularization and decidualization (19), the improvement of endometrial maturation (13), and the promotion of endometrial gene expression that may lead to a better synchrony between the embryo and the endometrium (20).

2 Hypothesis, objective and aim of the clinical trial

2.1 Study hypothesis [level 2]

Patients that receive ES during the cycle preceding the embryo transfer, have increased endometrial receptivity and thus higher clinical pregnancy rates (CPR).

2.1.1 Operative hypothesis [level 3]

Null hypothesis: Clinical pregnancy rate in patients who have undergone an ES in the previous cycle is not different from those who have not received it.

Alternative hypothesis: Patients who receive an ES in the previous cycle have a 15% higher CPR than those who have not received it.

2.2 Main outcome [level 2]

The main objective of the ENDOSCRATCH trial is to determine if there are differences in pregnancy rates in egg donor IVF treatments when comparing patients receiving an ES before endometrial preparation for embryo transfer and those who will not receive any intervention. CPR will be determined by the visualization of a intrauterine gestational sac via vaginal ultrasound at approximately 6 weeks pregnancy.

2.2.1 Secondary outcomes [level 3]

2.2.1.1 Efficacy secondary outcomes [level 4]

1. Secondary endpoints are biochemical pregnancy rate (BPR), ongoing pregnancy rate (OPR), implantation rate (IR), miscarriage rate (MR), live birth rate (LBR) and cumulative pregnancy rate (CumPR).
 - a. Biochemical pregnancy rate will be determined by the ratio between the number of patients with blood β -hCG levels over 10 mUI/ml 12 days after

- 111 the embryo transfer and the number of embryo transfers.
- 112 b. Ongoing pregnancy will be assessed via ultrasound beyond 12 weeks of
- 113 pregnancy.
- 114 c. Implantation rate will be determined by the ratio between the number of
- 115 gestational sacs and the number of replaced embryos.
- 116 d. Miscarriage rate will be determined by the ratio between the number of
- 117 miscarriages and the total number of pregnancies.
- 118 Early miscarriage will be assessed if pregnancy stops before the 12th week
- 119 of pregnancy. Late miscarriage will be assessed if pregnancy stops
- 120 between the 12th and the 24th week of pregnancy.
- 121 e. Live birth rate will be determined by the ratio between babies born and the
- 122 number of embryo transfers.
- 123 f. Multiple pregnancy rate, determined by the ratio between the number of
- 124 multiple pregnancies and the total number of pregnancies.
- 125 g. CumPR will be evaluated 12 months after randomization for all patients.
- 126 2. To determine the possible beneficial effect of ES in egg donor IVF recipients
- 127 adjusted by: age, BMI, smoking habits.
- 128 3. To evaluate the CPR in both groups regarding the donor's age, male partner's age
- 129 and sperm quality, number of eggs obtained and fertilized, total number of
- 130 embryos obtained and quality and number of transferred embryos.
- 131 4. To determine the possible positive effect in specific treatment subgroups:
- 132 a. Previous implantation failures from IVF with own eggs
- 133 b. Previous implantation failures from IVF with donor eggs
- 134 c. Previous miscarriages
- 135 d. Previous live births
- 136 5. To evaluate the interference of ES with the endometrial preparation, in terms of:
- 137 a. Endometrial thickness
- 138 b. Stimulation duration
- 139 c. Dose of medication needed
- 140

2.2.1.2 Safety secondary outcomes [level 4]

1. Obstetric outcome analysis to evaluate the possible association of this technique with:
 - a. Early or late miscarriage
 - b. Placentation anomalies
 - c. Intrauterine growth restriction
 - d. Preterm birth
 - e. Premature membrane rupture
 - f. Gestational diabetes
 - g. Gestational hypertension
2. Secondary effects of ES:
 - a. Pain
 - b. Bleeding

3 Clinical trial design [level 1]

3.1 Clinical trial description [level 2]

This is a single-centre prospective RCT, fully conducted at ProcreaTec Fertility Clinic in Madrid, starting January 2017 to December 2019 to evaluate the effectiveness of an endometrial biopsy (scratching) before endometrial preparation, during the luteal phase of the previous cycle versus the conventional treatment protocol for egg donation IVF without endometrial biopsy.

3.1.1 Clinical trial summarized description [level 3]

1. Study: Interventional.
2. Assignment: Randomized.
3. Final classification: Efficacy study.
4. Intervention model: Parallel assignment.
5. Blinding: Not blinded.
6. Main outcome: The improvement in clinical pregnancy rate in egg donor IVF recipients.

3.2 Patient randomization [level 2]

Patients starting egg donor IVF cycles that fulfill inclusion criteria will be offered participation. If they agree they will be assigned to a treatment group according to the randomization chart, which will be obtained by a web-based randomization program using random blocks (randomization.com). Since patients in the study group will receive an intervention and those in the control group will not (no placebo intervention will be performed), blinding is not possible for patients nor for physicians. All patients will sign IC to be enrolled in the study.

1. Group A (176 patients): Intervention group. They will receive an ES during the luteal phase of the previous cycle to embryo transfer.
2. Group B (176 patients): Control group. They will undergo the conventional protocol for donor IVF treatment.

3.3 Sample size calculation [level 2]

The average CPR after embryo transfer in egg donor IVF cycles is 60% at our centre. Based on previous studies, where the difference in CPR for IVF cycles varied between 10 to 30%(2,4,5,12,14,21–23), we estimated that a 15% difference in CPR would be clinically relevant. According to that percentage, a total of 332 patients will be needed to detect a 15% difference between the two groups, with 80% statistical power and two-sided alpha of 0,05. Considering a 5% dropout rate, we will include 176 patients per study arm, 352 patients in total.

3.4 Interim analysis [level 2]

Non applicable.

3.5 Study setting [level 2]

We will collect data from patients undergoing an egg donor IVF cycle at ProcreaTec Fertility Centre in Madrid from the 13th of January 2017.

3.5.1 Eligible population [level 2]

Those patients undergoing an egg donor IVF treatment protocol will be eligible for this study.

3.5.2 Study population [level 2]

Study population will include those patients fulfilling the inclusion criteria who have accepted the study and signed the IC.

3.5.3 Inclusion and exclusion criteria

3.5.3.1 Inclusion criteria:

Patients will be included if they meet the following inclusion criteria:

- Age between 18 and 50 years
- Primary or secondary infertility
- Patients undergoing an IVF protocol with donor eggs
- Normal uterine cavity (transvaginal ultrasound scan)
- Patients with endometrial polyps can be included as long as polypectomy is performed at least two months before the treatment cycle

3.5.3.2 Exclusion criteria:

Patients will be excluded if:

- There is a severe male factor (less than 2 million sperms per ml)
- They have uterine anomalies such as uterine fibroids that impact the cavity, Mullerian malformations or severe adenomyosis
- They have unilateral or bilateral hydrosalpinx
- They have undergone a previous ES or hysteroscopy (at least one month before the randomization)
- Pre-implantation genetic testing cycles

3.6 Data collection

All study variables will be collected from patients included in the trial, from ProcreaTec clinical records, according to the information required in the data collection form (Annex I). Each doctor will include relevant information in the patients' clinical record and the principal investigator will be responsible for collecting and managing the information. Any adverse events will be reported by responsible doctors and managed by the principal investigator.

3.7 Ethical approval [level 2]

This study will be conducted after the authorization of the Ethical Committee of Princesa Hospital in Madrid (Registry nº 2934/12-01-2017). Clinical data will be treated confidentially following the Spanish data protection law (Spanish Organic Law 15/1999, 13thDec).

4 Statistical principles [level 1]

4.1 General considerations

Baseline characteristics of patients included will be analyzed as follows. Qualitative variables will be described using mean and standard deviation, non-normal variables will be summarized using median and 25% and 75% centiles. Qualitative variables will be described using frequency distribution.

The main outcome, CPR, and secondary outcomes, BPR, OPR, MR, IR, LBR and CumPR for each group will be analyzed with Chi-Squared test or Fisher's exact test. Efficacy of the treatment will be described as absolute and relative frequencies, together with the association strength by raw risk ratio (RR) with 95% confidence intervals. In addition, a general linear model, with a log link and binomial distribution, will be used to estimate the strength of association between primary and secondary outcomes adjusted by independent variables.

Results will be presented as RR and 95% confidence intervals. Statistical significance will be 0,05 (5% both sides α error) for all comparisons. Statistical analysis will be done using Stata 13 for Windows (StataCorp LP, Texas).

All analyses will be performed by the 9th version of SAS system.

4.2 Study population [level 2]

The main statistical analysis will be by intention to treat (ITT). Patients allocated to a determined treatment group will be followed and evaluated as members of that group, without regards to the fulfillment of the planned treatment.

4.3 Patient flow diagram [level 2]

Patient diagram flow will be detailed with the CONSORT standards (fig 1).




Figure 1. CONSORT patient flow diagram.

4.4 Outcome evaluation

4.4.1 Efficacy outcome

The primary study objective will be to evaluate if endometrial scratching improves CPR in patients undergoing egg-donation IVF versus patients without endometrial scratching. The primary study objective will be performed using the Chi-square statistic test at a 2-sided significance level of $\alpha = 0.05$. The intervention effect will be quantified using the risk ratio (RR) with the 95% confidence interval and p-value (**Error! No se encuentra el origen de la referencia.**). The RR estimation will be realized using a generalized regression model with log link and binary outcome (clinical pregnancy equal to yes or no). The intervention RR will be adjusted by the following pre-treatment variables (Tabl): age, BMI, smoker, previous live birth, previous biochemical miscarriage, previous miscarriage, number of previous failed cycles with own eggs. Also, a multivariate model with the treatment will be performed to predict the CPR (Table 4). Finally, a predictive model will be generated with the treatment and the previous covariables.

	Group A (n=XXX)	Group B (n=XXX)	RR (IC 95%)	p-value
Biochemical pregnancy	NN (X%)	NN (X%)	XX (XX;XX)	P=XX
Clinical pregnancy	NN (X%)	NN (X%)	XX (XX;XX)	P=XX
Ongoing pregnancy	NN (X%)	NN (X%)	XX (XX;XX)	P=XX
Early miscarriage	NN (X%)	NN (X%)	XX (XX;XX)	P=XX
Late miscarriage	NN (X%)	NN (X%)	XX (XX;XX)	P=XX
Abortion	NN (X%)	NN (X%)	XX (XX;XX)	P=XX
Live birth	NN (X%)	NN (X%)	XX (XX;XX)	P=XX
Multiple pregnancy	NN (X%)	NN (X%)	XX (XX;XX)	P=XX
Cumulative clinical pregnancy rate (12 months)	NN (X%)	NN (X%)	XX (XX;XX)	P=XX

278 *Table 1. Clinical outcome*

	GROUP A (n=XXX)	GROUP B (n=XXX)
Race		
Arabian	XX (X%)	XX (X%)
Asian	XX (X%)	XX (X%)
Caucasian	XX (X%)	XX (X%)
Mixed	XX (X%)	XX (X%)
Hispanic	XX (X%)	XX (X%)
Mulatto	XX (X%)	XX (X%)
Black	XX (X%)	XX (X%)
Age	XX (XX)	XX (XX)
BMI	XX (XX)	XX (XX)
Smoking habit	XX (X%)	XX (X%)
Previous live birth	XX (X%)	XX (X%)
Previous miscarriage	XX (X%)	XX (X%)
Previous biochemical miscarriage	XX (X%)	XX (X%)
Previous ectopic pregnancy	XX (X%)	XX (X%)
Previous abortion	XX (X%)	XX (X%)
Number of previous failed cycles with own eggs		
0	XX (X%)	XX (X%)
1	XX (X%)	XX (X%)
2	XX (X%)	XX (X%)
+2	XX (X%)	XX (X%)
Number of previous failed cycles with donor eggs		
0	XX (X%)	XX (X%)
1	XX (X%)	XX (X%)
2	XX (X%)	XX (X%)
+2	XX (X%)	XX (X%)

	Group A (n=XXX)	Group B (n=XXX)	p-value
Donor's age	NN (X%)	NN (X%)	P=XX
Male partner's age	NN (X%)	NN (X%)	P=XX
Fresh sperm	NN (X%)	NN (X%)	P=XX
Frozen sperm	NN (X%)	NN (X%)	P=XX
Donor sperm	NN (X%)	NN (X%)	P=XX
Total sperm count	NN (X%)	NN (X%)	P=XX
Number of MII obtained	NN (X%)	NN (X%)	P=XX
Number of fertilized eggs	NN (X%)	NN (X%)	P=XX
Total number of embryos obtained	NN (X%)	NN (X%)	P=XX
Number of replaced embryos	NN (X%)	NN (X%)	P=XX
Quality of replaced embryos			
HATCHED	NN (X%)	NN (X%)	P=XX
HATCHING	NN (X%)	NN (X%)	P=XX
EXPANDED	NN (X%)	NN (X%)	P=XX
EARLY BLASTOCYST	NN (X%)	NN (X%)	P=XX
MORULA	NN (X%)	NN (X%)	P=XX
CLEAVAGE STAGE	NN (X%)	NN (X%)	P=XX

Adjusted model		
	RR (CI 95%)	p-value
Group A vs Group B	XX (XX;XX)	P=XX
Age	XX (XX;XX)	P=XX
Group A vs Group B	XX (XX;XX)	P=XX
BMI	XX (XX;XX)	P=XX
Group A vs Group B	XX (XX;XX)	P=XX
Smoking habit	XX (XX;XX)	P=XX
Group A vs Group B	XX (XX;XX)	P=XX
Previous live births	XX (XX;XX)	P=XX
Group A vs Group B	XX (XX;XX)	P=XX
Previous biochemical miscarriage	XX (XX;XX)	P=XX
Group A vs Group B	XX (XX;XX)	P=XX
Previous miscarriage	XX (XX;XX)	P=XX
Group A vs Group B	XX (XX;XX)	P=XX
Previous IVF own-eggs	XX (XX;XX)	P=XX
Group A vs Group B	XX (XX;XX)	P=XX
Previous IVF donor eggs	XX (XX;XX)	P=XX

Table 4. Model to evaluate the CPR and LBR according with the treatment and adjusted by age, BMI, smoking habit, previous live birth, previous biochemical miscarriage, previous miscarriage, previous IVF own eggs or previous IVF donor eggs failures.

Multivariate model		
	RR (CI 95%)	p-value
Endometrial scratching vs no intervention	XX (XX;XX)	P=XX
Age	XX (XX;XX)	P=XX
BMI	XX (XX;XX)	P=XX
Smoking habit	XX (XX;XX)	P=XX
Previous live birth	XX (XX;XX)	P=XX
Previous biochemical pregnancies	XX (XX;XX)	P=XX
Previous miscarriages	XX (XX;XX)	P=XX
Previous failed own eggs IVF embryo transfers	XX (XX;XX)	P=XX
Previous failed egg-donor IVF embryo transfers	XX (XX;XX)	P=XX

Table 5. Multivariate model to evaluate the CPR with the treatment adjusted by covariables

The following subgroups for the CPR will be defined (Table 6): previous implantation failures with own eggs, previous implantation failures with donor eggs, previous

miscarriage, previous live births. In each group, the effect of endometrial scratching (RR) for the CPR will be estimated using a generalized lineal model with log link and binary outcome (CPR equal yes or no). The comparison between groups will be realized using interaction between treatment and subgroup in the generalized linear model.

	GROUP A		GROUP B		
	CPR	Total	CPR	Total	RR (IC 95%)
Previous implantation failures with own eggs (p=XX)					
Yes	XX (XX%)	XX	XX (XX%)	XX	XX (XX;XX)
No	XX (XX%)	XX	XX (XX%)	XX	XX (XX;XX)
Previous implantation failures with donor eggs (p=XX)					
Yes	XX (XX%)	XX	XX (XX%)	XX	XX (XX;XX)
No	XX (XX%)	XX	XX (XX%)	XX	XX (XX;XX)
Previous miscarriages (p=XX)					
Yes	XX (XX%)	XX	XX (XX%)	XX	XX (XX;XX)
No	XX (XX%)	XX	XX (XX%)	XX	XX (XX;XX)
Previous live births (p=XX)					
Yes	XX (XX%)	XX	XX (XX%)	XX	XX (XX;XX)
No	XX (XX%)	XX	XX (XX%)	XX	XX (XX;XX)

Table 6. Effect of endometrial scratching on CPR by subgroups

The effect of endometrial scratching on secondary study objectives, LBR and CumPR, will be evaluate using the Chi-square statistic test at a 2-sided significance level of alpha = 0.05. The intervention effect will be quantified using the risk ratio (RR) with the 95% confidence interval and p-value (**¡Error! No se encuentra el origen de la referencia.**). The RR estimation will be realized using a generalized regression model with log link and binary outcome (clinical pregnancy equal to yes or no). The intervention RR will be adjusted by the following pre-treatment variables (Tabl): age, BMI, smoker, previous live

birth, previous biochemical miscarriage, previous miscarriage, number of previous failed cycles with own eggs and donor eggs.

The effect of endometrial scratching on stimulation duration, endometrial thickness and total dose required will be evaluate using the Student's t statistic test at a 2-sided significance level of $\alpha = 0.05$. If the normality hypothesis is not rejected with the Shapiro-Wilk test, the objective variables will be presented using the mean and the standard deviation. If the normality hypothesis is rejected, the variables will be analyzed with no-parametric test of Mann-Whitney and summarized with the median and range interquartile, 25 and 75 percentile. The data will be described according to the Table 7.

	GROUP A (n=XXX)	GROUP B (n=XXX)	p-value
Stimulation duration	XX (XX)	XX (XX)	P=XX
Endometrial thickness	XX (XX)	XX (XX)	P=XX
Total dose required	XX (XX)	XX (XX)	P=XX

Table 7. Results of endometrial preparation

4.4.2 Safety outcomes

The effect of endometrial scratching on pregnancy complications: miscarriage, placentation anomalies, intrauterine growth restriction, preterm birth, premature membrane rupture, gestational diabetes and gestational hypertension, will be evaluated using the Chi-square statistic test at a 2-sided significance level of $\alpha = 0.05$. The intervention effect will be quantified for each safety outcome using the risk ratio (RR) with the 95% confidence interval and p-value (**Error! No se encuentra el origen de la referencia.**). The RR estimation will be realized using a generalized regression model with log link and binary outcome (yes or no). The intervention RR will be summarized according to Table 8.

	GROUP A (n=XXX)	GROUP B (n=XXX)	RR (IC 95%)	p-value
Miscarriage	XX (X%)	XX (X%)	XX (XX;XX)	P=XX
Placentation abnomalies	XX (X%)	XX (X%)	XX (XX;XX)	P=XX
Intrauterine growth restriction	XX (X%)	XX (X%)	XX (XX;XX)	P=XX
Preterm birth	XX (X%)	XX (X%)	XX (XX;XX)	P=XX
Premature membrane rupture	XX (X%)	XX (X%)	XX (XX;XX)	P=XX
Gestational diabetes	XX (X%)	XX (X%)	XX (XX;XX)	P=XX
Gestational hypertension				

Table 8. Obstetrical outcome.

The absolute and relative frequencies of the pain and bleeding will be summarized by the patient with endometrial scratching. The endometrial preparation time will be described with the median, range interquartile, 25 and 75 percentile.

4.5 Baseline characteristics

The baseline characteristics: demographics, clinical, laboratory and relationship with the treatments will be presented with the mean and standard deviation values or absolute and relative frequency according to the treatment. If the distribution of continuous variable is not normal (Kolmogorov-Smirnov test), the information will be summarized with median and range interquartile, 25 and 75 percentile.

4.6 Evaluation of lost or unknown data.

The outlier data will be described with the maximum and minimum values. The outlier values will be revised with the CRD and corrected it. As loss to follow-up is expected to be minimal (i.e. less than one percent missing data on primary and secondary outcomes), missing values will not be imputed.

4.7 Additional considerations

When the fifty percent of the cells have expected counts less than five, the chi-square tests will be replaced by Fisher's exact test.

337 The protocol or statistical analysis plan deviation will be described and justified in the
338 deviation documents.

339 5 GLOSSARY

- 340 BMI: body mass index
- 341 BPR: Biochemical pregnancy rate
- 342 CI: confidence interval
- 343 COS: Controlled ovarian stimulation
- 344 CPR: Clinical pregnancy rate
- 345 CumPR: Cumulative pregnancy rate
- 346 ES: Endometrial scratching
- 347 IC: Informed consent
- 348 CI: Confidence Interval
- 349 IR: implantation rate
- 350 IUI: Intrauterine insemination
- 351 IVF: In vitro Fertilization
- 352 LBR: Live birth rate
- 353 MR: miscarriage rate
- 354 OPR: ongoing pregnancy rate
- 355 OR: Odds ratio
- 356 RR: Risk ratio

357

358 6 REFERENCES

- 359 1. De Geyter C, Calhaz-Jorge C, Kupka MS, Wyns C, Mocanu E, Motrenko T, et al.
360 ART in Europe, 2014: results generated from European registries by ESHRE: The

- 361 European IVF-monitoring Consortium (EIM) for the European Society of Human
362 Reproduction and Embryology (ESHRE). *Hum Reprod Oxf Engl*. 2018
363 01;33(9):1586–601.
- 364 2. Barash A, Dekel N, Fieldust S, Segal I, Schechtman E, Granot I. Local injury to the
365 endometrium doubles the incidence of successful pregnancies in patients
366 undergoing in vitro fertilization. *Fertil Steril*. 2003 Jun;79(6):1317–22.
- 367 3. Raziel A, Schachter M, Strassburger D, Bern O, Ron-El R, Friedler S. Favorable
368 influence of local injury to the endometrium in intracytoplasmic sperm injection
369 patients with high-order implantation failure. *Fertil Steril*. 2007 Jan;87(1):198–201.
- 370 4. Zhou L, Li R, Wang R, Huang H, Zhong K. Local injury to the endometrium in
371 controlled ovarian hyperstimulation cycles improves implantation rates. *Fertil Steril*.
372 2008 May;89(5):1166–76.
- 373 5. Karimzadeh MA, Ayazi Rozbahani M, Tabibnejad N. Endometrial local injury
374 improves the pregnancy rate among recurrent implantation failure patients
375 undergoing in vitro fertilisation/intra cytoplasmic sperm injection: a randomised
376 clinical trial. *Aust N Z J Obstet Gynaecol*. 2009 Dec;49(6):677–80.
- 377 6. Frantz S, Parinaud J, Kret M, Rocher-Escriva G, Papaxanthos-Roche A, Creux H,
378 et al. Decrease in pregnancy rate after endometrial scratch in women undergoing a
379 first or second in vitro fertilization. A multicenter randomized controlled trial. *Hum*
380 *Reprod Oxf Engl*. 2019 Jan 1;34(1):92–9.
- 381 7. Lensen S, Osavlyuk D, Armstrong S, Stadelmann C, Hennes A, Napier E, et al. A
382 Randomized Trial of Endometrial Scratching before In Vitro Fertilization. *N Engl J*
383 *Med*. 2019 24;380(4):325–34.
- 384 8. Yeung TWY, Chai J, Li RHW, Lee VCY, Ho PC, Ng EHY. The effect of
385 endometrial injury on ongoing pregnancy rate in unselected subfertile women
386 undergoing in vitro fertilization: a randomized controlled trial. *Hum Reprod Oxf*
387 *Engl*. 2014 Nov;29(11):2474–81.
- 388 9. Eskew AM, Reschke LD, Woolfolk C, Schulte MB, Boots CE, Broughton DE, et al.
389 Effect of endometrial mechanical stimulation in an unselected population
390 undergoing in vitro fertilization: fertility analysis of a double-blind randomized
391 controlled trial. *J Assist Reprod Genet*. 2018 Nov 5;
- 392 10. Baum M, Yerushalmi GM, Maman E, Kedem A, Machtinger R, Hourvitz A, et al.
393 Does local injury to the endometrium before IVF cycle really affect treatment
394 outcome? Results of a randomized placebo controlled trial. *Gynecol Endocrinol Off*
395 *J Int Soc Gynecol Endocrinol*. 2012 Dec;28(12):933–6.
- 396 11. Gibreel A, El-Adawi N, Elgindy E, Al-Inany H, Allakany N, Tournaye H.
397 Endometrial scratching for women with previous IVF failure undergoing IVF
398 treatment. *Gynecol Endocrinol Off J Int Soc Gynecol Endocrinol*. 2015
399 Apr;31(4):313–6.
- 400 12. Nastri C, Ferriani R, Raine-Fenning N, Martins W. Endometrial scratching
401 performed in the non-transfer cycle and outcome of assisted reproduction: a

- 402 randomized controlled trial. *Ultrasound Obstet Gynecol.* 2013;42(4):375-382.
- 403 13. Nastri CO, Lensen S, Polanski L, Raine-Fenning N, Farquhar CM, Martins WP.
404 Endometrial injury and reproductive outcomes: there's more to this story than meets
405 the horse's blind eye. *Hum Reprod Oxf Engl.* 2015 Mar;30(3):749.
- 406 14. Potdar N, Gelbaya T, Nardo LG. Endometrial injury to overcome recurrent embryo
407 implantation failure: a systematic review and meta-analysis. *Reprod Biomed Online.*
408 2012 Dec;25(6):561–71.
- 409 15. Nastri CO, Lensen SF, Gibreel A, Raine-Fenning N, Ferriani RA, Bhattacharya S,
410 et al. Endometrial injury in women undergoing assisted reproductive techniques.
411 *Cochrane Database Syst Rev* [Internet]. 2015 [cited 2019 Jun 7];(3). Available from:
412 [https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD009517.pub3/full](https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD009517.pub3/full?highlightAbstract=reproduct%7Cwoman%7Cin%7Cwithdrawn%7Cundergoing%7Cinjury%7Creproductive%7Ctechnique%7Cassisted%7Cendometrial%7Ctechniques%7Cendometri%7Creproductiv%7Cassist%7Cinjury%7Cundergo%7Cwomen)
413 [?highlightAbstract=reproduct%7Cwoman%7Cin%7Cwithdrawn%7Cundergoing%](https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD009517.pub3/full?highlightAbstract=reproduct%7Cwoman%7Cin%7Cwithdrawn%7Cundergoing%7Cinjury%7Creproductive%7Ctechnique%7Cassisted%7Cendometrial%7Ctechniques%7Cendometri%7Creproductiv%7Cassist%7Cinjury%7Cundergo%7Cwomen)
414 [7Cinjury%7Creproductive%7Ctechnique%7Cassisted%7Cendometrial%7Ctechniqu](https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD009517.pub3/full?highlightAbstract=reproduct%7Cwoman%7Cin%7Cwithdrawn%7Cundergoing%7Cinjury%7Creproductive%7Ctechnique%7Cassisted%7Cendometrial%7Ctechniques%7Cendometri%7Creproductiv%7Cassist%7Cinjury%7Cundergo%7Cwomen)
415 [es%7Cendometri%7Creproductiv%7Cassist%7Cinjury%7Cundergo%7Cwomen](https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD009517.pub3/full?highlightAbstract=reproduct%7Cwoman%7Cin%7Cwithdrawn%7Cundergoing%7Cinjury%7Creproductive%7Ctechnique%7Cassisted%7Cendometrial%7Ctechniques%7Cendometri%7Creproductiv%7Cassist%7Cinjury%7Cundergo%7Cwomen)
- 416 16. Dain L, Ojha K, Bider D, Levron J, Zinchenko V, Walster S, et al. Effect of local
417 endometrial injury on pregnancy outcomes in ovum donation cycles. *Fertil Steril.*
418 2014 Oct;102(4):1048–54.
- 419 17. Coughlan C, Ledger W, Wang Q, Liu F, Demirel A, Gurgan T, et al. Recurrent
420 implantation failure: definition and management. *Reprod Biomed Online.* 2014
421 Jan;28(1):14–38.
- 422 18. Cakmak H, Taylor HS. Implantation failure: molecular mechanisms and clinical
423 treatment. *Hum Reprod Update.* 2011 Apr;17(2):242–53.
- 424 19. Gnainsky Y, Granot I, Aldo PB, Barash A, Or Y, Schechtman E, et al. Local injury
425 of the endometrium induces an inflammatory response that promotes successful
426 implantation. *Fertil Steril.* 2010 Nov;94(6):2030–6.
- 427 20. Kalma Y, Granot I, Gnainsky Y, Or Y, Czernobilsky B, Dekel N, et al. Endometrial
428 biopsy-induced gene modulation: first evidence for the expression of bladder-
429 transmembranal uroplakin Ib in human endometrium. *Fertil Steril.* 2009
430 Apr;91(4):1042–9, 1049.e1-9.
- 431 21. Narvekar SA, Gupta N, Shetty N, Kottur A, Srinivas M, Rao KA. Does local
432 endometrial injury in the nontransfer cycle improve the IVF-ET outcome in the
433 subsequent cycle in patients with previous unsuccessful IVF? A randomized
434 controlled pilot study. *J Hum Reprod Sci.* 2010 Jan;3(1):15–9.
- 435 22. Seval MM, Şükür YE, Özmen B, Kan Ö, Sönmezer M, Berker B, et al. Does adding
436 endometrial scratching to diagnostic hysteroscopy improve pregnancy rates in
437 women with recurrent in-vitro fertilization failure? *Gynecol Endocrinol Off J Int*
438 *Soc Gynecol Endocrinol.* 2016 Dec;32(12):957–60.
- 439 23. Maged AM, Rashwan H, AbdelAziz S, Ramadan W, Mostafa WAI, Metwally AA,
440 et al. Randomized controlled trial of the effect of endometrial injury on implantation
441 and clinical pregnancy rates during the first ICSI cycle. *Int J Gynaecol Obstet Off*
442 *Organ Int Fed Gynaecol Obstet.* 2018 Feb;140(2):211–6.

